

JOURNAL OF ALGEBRA 22, 461–479 (1972)

Integral Relative Grothendieck Rings*

WILLIAM H. GUSTAFSON

*Department of Mathematics, Indiana University, Bloomington, Indiana 47401**Communicated by J. A. Green*

Received October 13, 1970

1. INTRODUCTION

Let R be a Dedekind ring and G a finite group. An RG -lattice is an RG -module which is finitely generated and projective as an R -module. Let H be a fixed subgroup of G . Form the free abelian group X on the symbols $[M]$, where M ranges over a full set of representatives of the isomorphism classes of RG -lattices. Denote by Y the subgroup of X generated by all elements $[M] - [M'] - [M'']$ such that there is an RG -exact sequence

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0,$$

which is split when viewed as a sequence of RH -lattices. The quotient group X/Y will be denoted by $a_R(G, H)$. The element $[M] + Y \in a_R(G, H)$ will also be denoted by $[M]$. If M, N are RG -lattices, define $M \otimes N$ to be the RG -lattice obtained by letting G act on $M \otimes_R N$ through the formula

$$g \cdot (m \otimes n) = gm \otimes gn, \quad g \in G, \quad m \in M, \quad n \in N.$$

Then putting $[M] \cdot [N] = [M \otimes N]$, we give $a_R(G, H)$ the structure of a commutative ring with unit $[R]$, where G acts trivially on R . $a_R(G, H)$ is the *Grothendieck ring of RG -lattices relative to RH -split sequences*. When $G = H$, we write $a_R(G)$ instead of $a_R(G, G)$, for brevity.

In the event that R is a field of characteristic $p > 0$, $a_R(G, H)$ has been studied extensively by Lam, Reiner, Wigner, Dress, and Conlon (See [2]–[4], [6]–[12]). In this paper, we calculate some examples in the case where R is not a field. Our examples show that the methods of the “modular” case do not generalize to the “integral” case.

* This work supported by an NSF Traineeship at the University of Illinois. The author is indebted to Professor Irving Reiner for his advice and encouragement. He is also indebted to Professor J. A. Green for his useful comments on certain matters of style.

2. GROUPS OF PRIME ORDER

Let p be a rational prime. Denote by R the ring of p -adic integers in the field K of p -adic numbers, and let $H = \langle h \rangle$ denote a cyclic group of order p . Denote by A the RH -lattice R with trivial action of H . Let ζ be a primitive p -th root of unity over K , and denote by B the RH -lattice $R[\zeta]$, where $h \cdot x = \zeta \cdot x$ for $x \in R[\zeta]$.

(2.1) THEOREM (Heller and Reiner [5]). *A and B are the only irreducible RH -lattices. The indecomposable RH -lattices are A , B , and RH . There are nonsplit RH -exact sequences*

$$\begin{aligned} 0 \rightarrow A \rightarrow RH \rightarrow B \rightarrow 0, \\ 0 \rightarrow B \rightarrow RH \rightarrow A \rightarrow 0. \end{aligned}$$

Further,

$$\text{ext}_{RH}^1(A, B) \cong \text{ext}_{RH}^1(B, A) \cong A/pA \cong \mathbb{Z}/p\mathbb{Z},$$

and if

$$\begin{aligned} 0 \rightarrow A \rightarrow X \rightarrow B \rightarrow 0 \\ 0 \rightarrow B \rightarrow Y \rightarrow A \rightarrow 0 \end{aligned}$$

are nonsplit exact sequences, then $X \cong Y \cong RH$.

Now let t, u be positive integers. Denote by $A^{(u)}$ the RH -lattice $A \oplus \cdots \oplus A$ (u copies), etc. Then we have

$$(2.2) \quad \text{ext}_{RH}^1(B^{(t)}, A^{(u)}) \cong (A/pA)_{t \times u},$$

where $(A/pA)_{t \times u}$ denotes the module of all $t \times u$ matrices over A/pA . This isomorphism may be described explicitly as follows: Let

$$\begin{aligned} \Phi &= 1 + h + h^2 + \cdots + h^{p-1}. \text{ Write} \\ B^{(t)} &= B \cdot b_1 \oplus \cdots \oplus B \cdot b_t \\ A^{(u)} &= A \cdot a_1 \oplus \cdots \oplus A \cdot a_u. \end{aligned}$$

Let $RH \cdot x_1 \oplus \cdots \oplus RH \cdot x_t$ be a free RH -module with basis x_1, \dots, x_t . Define $\tau: \sum RH \cdot x_i \rightarrow \sum B \cdot b_i$ by $\tau(x_i) = b_i$ for $i = 1, \dots, t$. Then $\ker \tau = \sum RH \cdot \Phi \cdot x_i$, so there is an exact sequence

$$0 \longrightarrow \sum RH \cdot \Phi \cdot x_i \longrightarrow \sum RH \cdot x_i \xrightarrow{\tau} \sum B \cdot b_i \longrightarrow 0.$$

Hence there is an exact sequence

$$\begin{aligned} \operatorname{hom}_{RH} \left(\sum RH \cdot x_i, \sum A \cdot a_j \right) &\xrightarrow{\tau^*} \operatorname{hom}_{RH} \left(\sum RH \cdot \Phi \cdot x_i, \sum A \cdot a_j \right) \\ &\rightarrow \operatorname{ext}_{RH}^1 \left(\sum B \cdot b_i, \sum A \cdot a_j \right) \rightarrow \operatorname{ext}_{RH}^1 \left(\sum RH \cdot x_i, \sum A \cdot a_j \right). \end{aligned}$$

Since $\sum RH \cdot x_j$ is RH -free, the last term is zero; therefore

$$\operatorname{ext}_{RH}^1 \left(\sum B \cdot b_i, \sum A \cdot a_j \right) \cong \operatorname{coker} \tau^*.$$

If $F: \sum RH \cdot \Phi \cdot x_i \rightarrow \sum A \cdot a_j$, then the matrix corresponding to $F + \operatorname{Im} \tau^*$ under the isomorphism (2.2) is obtained by writing $F(\Phi x_i) = \sum_j \alpha_{ij} a_j$, $\alpha_{ij} \in A$, and reducing the matrix (α_{ij}) modulo pA . The extension corresponding to $F + \operatorname{Im} \tau^*$ is the lower row of

$$\begin{array}{ccccccc} 0 & \longrightarrow & \sum RH \cdot \Phi x_i & \longrightarrow & \sum RH \cdot x_i & \longrightarrow & \sum B \cdot b_i \longrightarrow 0 \\ & & \downarrow F & & \downarrow & & \downarrow id \\ 0 & \longrightarrow & \sum A \cdot a_j & \longrightarrow & M & \longrightarrow & \sum B \cdot b_i \longrightarrow 0, \end{array}$$

where the left hand square is constructed as a pushout diagram. If r is the R/pR -rank of $(\alpha_{ij} + pA)$, then it follows from [5] that

$$M \cong RH^{(r)} \oplus A^{(u-r)} \oplus B^{(t-r)}.$$

(2.3) LEMMA [5]. *Let X be an RH -lattice. Then*

- (i) $\operatorname{ext}_{RH}^1(RH, X) = 0$
- (ii) $\operatorname{ext}_{RH}^1(X, RH) = 0$
- (iii) $\operatorname{ext}_{RH}^1(A, A) = 0$
- (iv) $\operatorname{ext}_{RH}^1(B, B) = 0$.

(2.4) LEMMA. *Suppose that*

$$E: 0 \longrightarrow A \xrightarrow{\alpha} M \longrightarrow M/A \longrightarrow 0$$

is a nonsplit short exact sequence of RH -lattices. Then there is an RH -sublattice X of M such that

- (i) $X \cong RH$
- (ii) $\alpha(A) \subseteq X$
- (iii) X is an RH -direct summand of M .

Proof. Write M/A as a direct sum of indecomposable lattices. Then there is a corresponding decomposition of $\text{ext}_{RH}^1(M/A, A)$. Since E represents a nonzero element of $\text{ext}_{RH}^1(M/A, A)$, it follows that for some indecomposable summand Y of M/A , the component of the class of E in $\text{ext}_{RH}^1(Y, A)$ is nonzero. By (2.3), we conclude immediately that $Y \cong B$. Since the projections in the decomposition of $\text{ext}_{RH}^1(M/A, A)$ are induced by the injections in the decomposition of M/A , we see that there is an RH -split exact sequence

$$0 \longrightarrow B \xrightarrow{i} M/A \longrightarrow N \longrightarrow 0$$

for some RH -lattice N such that $i^*(\text{class of } E) \neq 0$, where

$$i^* = \text{ext}_{RH}^1(i, A) : \text{ext}_{RH}^1(M/A, A) \rightarrow \text{ext}_{RH}^1(B, A).$$

It follows that any representative of $i^*(\text{class of } E)$ is nonsplit. Using the construction of i^* (see [13]), we see that there is, therefore, a diagram of RH -lattices

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \longrightarrow & X' & \longrightarrow & B \longrightarrow 0 \\ & & \downarrow \text{id} & & \downarrow \sigma & & \downarrow i \\ 0 & \longrightarrow & A & \xrightarrow{\alpha} & M & \longrightarrow & M/A \longrightarrow 0 \end{array}$$

in which the upper row is nonsplit and the right-hand square is a pullback. It is easy to verify that σ is monic and $\text{coker } \sigma \cong N$. Now, $X' \cong RH$ by (2.1), so by (2.3(ii)), $\sigma(X')$ is an RH -summand of M . Hence taking $X = \sigma(X')$, conditions (i)–(iii) are satisfied, and so the proof is complete.

We conclude this section with

(2.5) LEMMA. *Suppose a diagram*

$$\begin{array}{ccc} A & \xrightarrow{\alpha} & RH \\ \beta \downarrow & & \\ RH & & \end{array}$$

is given, where α, β are RH -monomorphisms with R -torsionfree cokernels. Then there are RH -homomorphisms $\mu, \theta: RH \rightarrow RH$ such that the diagram

$$\begin{array}{ccc} A & \xrightarrow{\alpha} & RH \\ \beta \downarrow & \nearrow \mu & \nearrow \theta \\ RH & & \end{array}$$

is commutative.

Proof. Straightforward.

3. CYCLIC p -GROUPS

In this section, G is a cyclic group of order p^n , and R is the ring of p -adic integers. Let g be a generator for G , and let $h = g^{p^{n-1}}$. Then $H = \langle h \rangle$ is the unique subgroup of order p of G . We will calculate the ring $a_R(G, H)$. The methods used can also be used in the case $G = H' \times H$, where H' is cyclic of order p^{n-1} , and H is again cyclic of order p . We omit the details of this case. The notation for RH -lattices established in §2 will be carried over without further mention.

It is not difficult to see that the irreducible RG -lattices are X_0, \dots, X_n , where for $i = 0, 1, \dots, n$, ζ_i is a primitive p^i -th root of unity, and $X_i = R[\zeta_i]$ with $g \cdot x = \zeta_i x$ for $x \in R[\zeta_i]$.

For any RH -lattice N , denote by N^G the induced lattice $RG \otimes_{RH} N$. Then the exact sequence

$$0 \rightarrow A \rightarrow RH \rightarrow B \rightarrow 0$$

gives rise to an exact sequence

$$(3.1) \quad 0 \rightarrow A^G \rightarrow RG \rightarrow B^G \rightarrow 0.$$

Further, the image of A in RH is just $RH \cdot \Phi$, so it follows readily that the image of A^G in RG is $RG \cdot \Phi$.

(3.2) PROPOSITION. *Let N be any H -trivial RG -lattice. Then*

$$\text{ext}_{RG}^1(B^G, N) \cong N/pN$$

as RG -modules.

Proof. The exact sequence (3.1) induces an exact sequence

$$\begin{aligned} \text{hom}_{RG}(RG, N) &\xrightarrow{\tau^*} \text{hom}_{RG}(RG \cdot \Phi, N) \longrightarrow \text{ext}_{RG}^1(B^G, N) \\ &\longrightarrow \text{ext}_{RG}^1(RG, N). \end{aligned}$$

The last term is zero, hence $\text{ext}_{RG}^1(B^G, N) \cong \text{coker } \tau^*$. Now, an RG -homomorphism $F: RG \cdot \Phi \rightarrow N$ is completely determined by $F(\Phi)$. Since the annihilator ideal of Φ is $RG \cdot (h - 1)$, and $(h - 1)N = 0$ by hypothesis, $F(\Phi)$ may be given any value in N . Thus $\text{hom}_{RG}(RG \cdot \Phi, N) \cong N$. The map τ^* is given by restriction, hence if $F: RG \rightarrow N$, then $\tau^*(F)(\Phi) = F(\Phi) = \Phi \cdot F(1)$. Now $F(1)$ may take any value in N , so in the isomorphism $\text{hom}_{RG}(RG \cdot \Phi, N) \cong N$, the image of τ^* is identified with $\Phi \cdot N$. But h acts trivially on N , so $\Phi \cdot N = p \cdot N$ and hence

$$\text{ext}_{RG}^1(B^G, N) \cong N/pN,$$

as claimed.

The above proof shows that for any element $n \in N$, we may construct an extension $(B^G, N; n)$ of B^G by N . Namely, we take $F(\Phi) = n$, and we construct the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & RG \cdot \Phi & \xrightarrow{\tau} & RG & \longrightarrow & B^G \longrightarrow 0 \\ & & F \downarrow & & \downarrow & & \downarrow id \\ 0 & \longrightarrow & N & \longrightarrow & M & \longrightarrow & B^G \longrightarrow 0, \end{array}$$

where the left-hand square is a pushout. We put $(B^G, N; n) = M$. Furthermore, the two extensions $(B^G, N; n)$ and $(B^G, N; n')$ are equivalent if and only if $n - n' \in pN$. It is easy to prove

(3.3) LEMMA. *Let α be an RG -automorphism of N . Then $(B^G, N; n) \cong (B^G, N; \alpha(n))$.*

Using the fact that $B^G \cong RG/RG \cdot \Phi$, it is easy to deduce:

(3.4) LEMMA. $X_n \cong B^G$. $(X_n)_H \cong B^{(p^{n-1})}$, where $(X_n)_H$ denotes X_n viewed as an RH -lattice by restriction of operators.

From the remarks above, we deduce that we may give extensions of X_n by X_0 by giving an element $x_0 \in X_0$. Let us use (3.3) to show

(3.5) LEMMA. *Up to isomorphism (of RG -lattices), there is only one non-split extension of X_n by X_0 .*

Proof. Let $x_0 \in X_0$ and consider $(X_n, X_0; x_0)$. Since R is a discrete valuation ring and $X_0 = R$, we may write $x_0 = p^k \cdot u$, where k is a nonnegative integer and u is a unit. The map $X_0 \rightarrow X_0$, given by multiplication by u^{-1} , is an RG -automorphism of X_0 , so by (3.3), we have

$$(X_n, X_0; x_0) \cong (X_n, X_0; p^k).$$

If $k \geq 1$, then $p^k \in pX_0$; so, $(X_n, X_0; p^k) \cong (X_n, X_0; 0) = X_n \oplus X_0$. Hence up to isomorphism, the only nonsplit extension of X_n by X_0 is $(X_n, X_0; 1)$.

Henceforth, $(X_n, X_0; 1)$ will be denoted by (X_n, X_0) .

Let $\Gamma = \langle \gamma \rangle$ be a cyclic group of order p^{n-1} . We identify Γ with G/H ; thus we have an epimorphism $\varphi: G \rightarrow \Gamma$ with kernel H . φ induces an epimorphism $RG \rightarrow R\Gamma$ through which we may regard $R\Gamma$ and its ideals as RG -lattices. In particular, let E denote $R\Gamma$ viewed as RG -lattice, and V the RG -sublattice $(\gamma - 1)R\Gamma$ of E . Observe that V is the augmentation ideal of $R\Gamma$.

Since $H = \ker \varphi$, E and V are H -trivial. Further, by [5, Theorem 1.7], E is a local ring with maximal ideal $pE + V$, and E/pE is a local principal ideal ring with maximal ideal V/pV . From this, it follows easily that any element of E is congruent modulo pE to an element of the form $(\gamma - 1)^k u$, with u a unit. Hence any extension of X_n by E is equivalent to an extension of the form $(X_n, E; (\gamma - 1)^k u)$. Now, if $k \geq p^{n-1}$, then $(\gamma - 1)^k \in pE$, whence $(X_n, E; (\gamma - 1)^k u) \cong X_n \oplus E$. On the other hand, if $k < p^{n-1}$, then by (3.3), we deduce

$$(X_n, E; (\gamma - 1)^k u) \cong (X_n, E; (\gamma - 1)^k).$$

Hence we have shown

(3.6) LEMMA. *All isomorphism types of RG -lattices which are extensions of B^G by E occur in the list $(X_n, E; 1), (X_n, E; (\gamma - 1)), \dots, (X_n, E; (\gamma - 1)^{p^{n-1}})$.*

Henceforth, $(X_n, E; (\gamma - 1)^k)$ will be denoted by $(X_n, E)_k$, for $0 \leq k \leq p^{n-1}$. Similarly, we denote $(X_n, V; (\gamma - 1)^k)$ by $(X_n, V)_k$, for $0 \leq k \leq p^{n-1}$.

Now we do some preliminary calculations in $a_R(G, H)$. Observe that there is an RG -exact sequence

$$(3.7) \quad 0 \longrightarrow V \xrightarrow{i} E \xrightarrow{\epsilon} X_0 \longrightarrow 0,$$

where ϵ is the augmentation map and i is the inclusion. This is an RH -split sequence. It is not difficult to verify that (3.7) induces an RH -split sequence

$$0 \rightarrow (X_n, V)_r \rightarrow (X_n, E)_r \rightarrow X_0 \rightarrow 0,$$

whence

$$(3.8) \quad [(X_n, E)_r] = [(X_n, V)_r] + [X_0] \quad \text{in } a_R(G, H), \quad \text{for } r = 0, 1, \dots, p^{n-1}.$$

We also have an RG -epimorphism $\sigma: E \rightarrow V$ given by $\sigma(e) = (\gamma - 1)e$, $e \in E$. Form the pushout diagram

$$\begin{array}{ccc} 0 \longrightarrow E & \longrightarrow & (X_n, E)_r \\ \sigma \downarrow & & \downarrow \delta \\ & V & \xrightarrow{\lambda} M. \end{array}$$

Then λ is monic, and $\ker \delta \cong \ker \sigma \cong X_0$. Let $F: RG \cdot \Phi \rightarrow E$ be given by $F(\Phi) = (\gamma - 1)^r$. Then the diagram

$$\begin{array}{ccc} RG \cdot \Phi & \xrightarrow{\tau} & RG \\ \sigma \circ F \downarrow & & \downarrow \\ & V & \xrightarrow{\lambda} M \end{array}$$

is a pushout, where $RG \rightarrow M$ is the composite of δ with the canonical map $RG \rightarrow (X_n, E)_r$. Hence $M \cong (X_n, V; \sigma F(\Phi))$. However, $\sigma F(\Phi) = (\gamma - 1)^{r+1}$, so $M \cong (X_n, V)_{r+1}$ by the uniqueness of pushouts. Thus, there is an RG -exact sequence

$$0 \longrightarrow X_0 \xrightarrow{\alpha} (X_n, E)_r \longrightarrow (X_n, V)_{r+1} \longrightarrow 0. \quad (3.9)$$

We will show below that (3.9) is not RH -split.

Suppose now that M is any extension of an H -trivial RG -lattice N , by $X_n(\cong B^G)$. We may construct a commutative diagram with RG -exact rows

$$\begin{array}{ccccccc} 0 & \rightarrow & RG \cdot \Phi & \rightarrow & RG & \rightarrow & X_n \rightarrow 0 \\ & & \downarrow F & & \downarrow & & \downarrow id \\ 0 & \longrightarrow & N & \longrightarrow & M & \rightarrow & X_n \rightarrow 0, \end{array}$$

by the remarks following the proof of Prop. (3.2). In order to calculate the isomorphism class of the restricted lattice M_H , it follows from the discussion of the isomorphism (2.2) and the structure of $(X_n)_H$ that we must choose an R -basis a_1, \dots, a_u for N , write

$$F((g-1)^i \Phi) = \sum_{j=1}^u \alpha_{ij} a_j, \quad \text{with } \alpha_{ij} \in R(0 \leq i \leq p^{n-1} - 1),$$

and then calculate the R/pR -rank s of the matrix $(\alpha_{ij} + pR)$. Once this is done, we will have

$$M_H \cong RH^{(s)} \oplus A^{(u-s)} \oplus B^{(p^{n-1}-s)}.$$

Let us apply this procedure to the modules $M = (X_n, E)_r$, $(X_n, V)_r$ and (X_n, X_0) . The module E has R -basis $\{(\gamma - 1)^i : 0 \leq i \leq p^{n-1} - 1\}$, and $(X_n, E)_r$ is determined by the homomorphism $F: RG \cdot \Phi \rightarrow E$ given by $F(\Phi) = (\gamma - 1)^r$. Thus $F((g-1)^i \cdot \Phi) = (g-1)^i \cdot (\gamma - 1)^r = (\gamma - 1)^{r+i}$. Since $(\gamma - 1)^{p^{n-1}} \in pE$, the matrix to be considered is the $p^{n-1} \times p^{n-1}$ matrix

$$\begin{pmatrix} 000 & \cdots & 010 & \cdots & 0 \\ 000 & \cdots & 001 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdot \\ 000 & \cdots & 000 & \cdots & 1 \\ 000 & \cdots & 000 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdot \\ 000 & \cdots & 000 & \cdots & 0 \end{pmatrix}, \quad (3.10)$$

where the first nonzero entry of the first row occurs in the $(r + 1)^{\text{st}}$ column. Clearly the rank of this matrix is $p^{n-1} - r$, hence

(3.11) PROPOSITION. $((X_n, E)_r)_H \cong RH^{(p^{n-1}-r)} \oplus A^{(r)} \oplus B^{(r)}$, for $0 \leq r \leq p^{n-1}$.

Similarly, one establishes

(3.12) PROPOSITION. $((X_n, V)_r)_H \cong RH^{(p^{n-1}-r)} \oplus A^{(r-1)} \oplus B^{(r)}$, for $1 \leq r \leq p^{n-1}$.

(3.13) COROLLARY. *The sequence (3.9) is not RH-split, for $0 \leq r \leq p^{n-1} - 1$.*

Proof. By (3.11), (3.12) and the Krull-Schmidt Theorem for RH-lattices [15].

Using the method above, one may also establish

(3.14) PROPOSITION. $(X_n, X_0)_H \cong RH \oplus B^{(p^{n-1}-1)}$;

and hence

(3.15) COROLLARY. *The sequence*

$$0 \longrightarrow X_0 \xrightarrow{\beta} (X_n, X_0) \longrightarrow X_n \longrightarrow 0$$

is not RH-split.

Now, by the argument in [8, Lemma (3.1)], one may show

(3.16) LEMMA. *Let R be a Dedekind ring, $G \supseteq H$ finite groups. Let M, N, L be RG -lattices and suppose that there exists a commutative diagram*

$$\begin{array}{ccc} M & \xrightarrow{\alpha} & N \\ \beta \downarrow & \nearrow \theta & \\ & \mu \nearrow & \\ L & & \end{array}$$

such that

- (i) α, β are RG -monomorphisms,
- (ii) $\text{coker } \alpha, \text{coker } \beta$ are RG -lattices,
- (iii) μ, θ are RH -homomorphisms.

Then

$$[N] - [\text{coker } \alpha] = [L] - [\text{coker } \beta] \quad \text{in } a_R(G, H).$$

(3.17) COROLLARY. Let R, G, H, M, N, L, α and β be as in (3.16). Suppose there are RH -submodules X of N_H , Y of L_H such that

- (i) X is an RH -summand of N_H ,
- (ii) Y is an RH -summand of L_H ,
- (iii) $\text{Im } \alpha \subseteq X, \text{Im } \beta \subseteq Y$,
- (iv) There are RH -homomorphisms $\mu_1 : X \rightarrow Y, \theta_1 : Y \rightarrow X$ with $\mu_1 \alpha = \beta, \theta_1 \beta = \alpha$.

Then

$$[N] - [\text{coker } \alpha] = [L] - [\text{coker } \beta] \quad \text{in } a_R(G, H).$$

Now let $0 \leq r \leq p^{n-1} - 1$. By (2.4), (2.5), (3.13), and (3.15), we may find a commutative diagram

$$\begin{array}{ccc} X_0 & \xrightarrow{\alpha} & Y \subseteq (X_n, E)_r \\ \beta \downarrow & \nearrow \theta_1 & \\ & & Z \subseteq (X_n, X_0), \\ & \nwarrow \mu_1 & \end{array}$$

where θ_1, μ_1 are RH -isomorphisms, $Z \cong Y \cong RH$, Z is an RH -summand of (X_n, X_0) , and Y is an RH -summand of $(X_n, E)_r$. By (3.17), we conclude

(3.18) PROPOSITION. For $0 \leq r \leq p^{n-1} - 1$,

$$[(X_n, X_0)] - [X_n] = [(X_n, E)_r] - [(X_n, V)_{r+1}] \quad \text{in } a_R(G, H).$$

(3.19) COROLLARY. $[(X_n, E)_{p^{n-1}-1}] = [(X_n, X_0)] + [V]$ in $a_R(G, H)$.

Comparing the expressions for $[(X_n, V)_r]$ obtained from (3.8) and (3.18), we obtain

(3.20) COROLLARY. $[(X_n, E)_{r-1}] = [(X_n, E)_r] - [X_0] - [X_n] + [(X_n, X_0)]$ in $a_R(G, H)$, for $1 \leq r \leq p^{n-1}$.

Using (3.20) repeatedly, along with (3.19), we obtain

(3.21) PROPOSITION. The subgroup of $a_R(G, H)$ generated by $[X_0], \dots, [X_n], [(X_n, E)_0], \dots, [(X_n, E)_{p^{n-1}}]$ is also generated by $[X_0], \dots, [X_n], [(X_n, X_0)]$.

Proof. We need only note that by (2.3(iii)), we have

$$[V] = [X_1] + \dots + [X_{n-1}] \quad \text{in } a_R(G, H).$$

Now we have two tasks remaining. Namely, we will show that $a_R(G, H)$ is generated by all $[X_i]$ and all $[(X_n, E)_k]$. Then it will follow from (3.21) that $a_R(G, H)$ is generated by $[X_0], \dots, [X_n], [(X_n, X_0)]$. Finally, we will show that this latter set of generators is independent.

We have seen that each irreducible RG -lattice is either H -trivial, or has the property that $Y_H \cong B^{(p^{n-1})}$. Let us say that an RG -lattice M is of *type 1* if $M_H \cong A^{(u)}$ for some u , and of *type 2* if $M_H \cong B^{(t)}$ for some t . We have just observed that each irreducible lattice belongs to one of the types. There exist lattices which are of neither type.

(3.22) LEMMA. *If M is of type 1 or of type 2, then in $a_R(G, H)$, $[M]$ equals the sum of its composition factors.*

Proof. Clear from (2.3).

(3.23) PROPOSITION. *Let U be the subgroup of $a_R(G, H)$ generated by all $[X_i]$ and all $[(X_n, E)_k]$. Then $U = a_R(G, H)$.*

Proof. Let M be an RG -lattice; we show that $[M] \in U$ by induction on the R -rank of M . Clearly if $\text{rank } M = 0$ or if M is of type 1 or 2, then $[M] \in U$.

Hence we may assume that M is of neither type. We may then find an exact sequence

$$(3.24) \quad 0 \longrightarrow I \xrightarrow{j} M \xrightarrow{k} N \longrightarrow 0$$

of RG -lattices, with I irreducible of type 1. Also, N admits an R -pure irreducible RG -lattice of type 2. If (3.24) is RH -split, then $[M] = [I] + [N]$ belongs to U by the induction hypothesis. If (3.24) is not RH -split, we must use a much more difficult argument. Let $N' = \{n \in N \mid \Phi n = 0\}$ be the maximal type 2 sublattice of N . Then there is an exact sequence

$$0 \longrightarrow N' \xrightarrow{\iota} N \longrightarrow W \longrightarrow 0,$$

where W is an H -trivial RG -lattice. We form the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & I & \xrightarrow{j'} & P & \longrightarrow & N' \longrightarrow 0 \\ & & \text{id} \downarrow & & \downarrow \sigma & & \downarrow i \\ 0 & \longrightarrow & I & \longrightarrow & M & \longrightarrow & N \longrightarrow 0, \end{array}$$

where the right-hand square is a pullback. Then σ is monic, and $\text{coker } \sigma \cong W$. Fix an isomorphism

$$\lambda: P_H \cong RH^{(u)} \oplus A^{(v)} \oplus B^{(w)}.$$

Since $\text{Im } j'$ is H -trivial, we have $\lambda(\text{Im } j') \subseteq RH^{(u)} \oplus A^{(v)}$. Thus $\text{Im } j'$ is contained in an RH -summand Y of P , where $Y \cong RH^{(u)} \oplus A^{(v)}$. Let $f: P_H \rightarrow Y$ be the canonical RH -projection onto the summand Y . Form the diagram of RH -modules

$$\begin{array}{ccccccc} 0 & \longrightarrow & P_H & \xrightarrow{\sigma} & M_H & \longrightarrow & W_H \longrightarrow 0 \\ & & f \downarrow & & \downarrow f' & & \downarrow id \\ 0 & \longrightarrow & Y & \xrightarrow{\sigma'} & M' & \longrightarrow & W_H \longrightarrow 0, \end{array}$$

where the left-hand square is a pushout diagram. Since $Y \cong RH^{(u)} \oplus A^{(v)}$ and W is H -trivial, it follows from (2.3) that there is an RH -homomorphism $\mu: M' \rightarrow Y$ with $\mu\sigma' = \text{Id}_Y$. Let $\theta: Y \rightarrow P_H$ be the inclusion, so that $f\theta = \text{Id}_Y$. Then $\mu f'\sigma\theta = \mu\sigma'f\theta = \text{Id}_Y \circ \text{Id}_Y = \text{Id}_Y$. Thus, $\sigma\theta$ is split by $\mu f'$; so $\sigma(Y)$ is an RH -summand of M_H . Now, $\sigma|_Y: Y \rightarrow \sigma(Y)$ has an inverse τ , and hence we have a commutative diagram

$$\begin{array}{ccc} I & \xrightarrow{j'} & Y \\ \downarrow & \swarrow \sigma & \nearrow \tau \\ \sigma(Y). & & \end{array}$$

Since Y is an RH -summand of P_H and $\sigma(Y)$ is an RH -summand of M_H , we may apply (3.17) to conclude

$$[M] = [P] - [N'] + [N] \quad \text{in } a_R(G, H).$$

Now, the R -ranks of N and N' are smaller than that of M , hence $[N']$, $[N] \in U$. Thus it suffices to show $[P] \in U$.

In order to deal with P , we must repeat the entire procedure with some slight variation. Thus, using P in place of M , and an R -pure, irreducible RG -sublattice J of type 2 in P in place of I , we work as before. We obtain

$$[P] = [P'] - [I] + [P/J] \quad \text{in } a_R(G, H),$$

where P' is such that there is a short exact sequence

$$0 \rightarrow J \rightarrow P' \rightarrow I \rightarrow 0.$$

Let $P'' = \{x \in P: (h-1)x = 0\}$. Then $P'' \cong I$ and $P'/P'' \cong J$. Thus, we have an exact sequence

$$0 \rightarrow I \rightarrow P' \rightarrow J \rightarrow 0.$$

Since E is the representation of G on the cosets of H , we have $E \cong A^G$. As A^G is the maximal type 1 sublattice of RG , and I is an R -pure type 1

sublattice of RG , we have an embedding $s: I \rightarrow E$ with R -pure image. Form the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & I & \longrightarrow & P' & \longrightarrow & J \longrightarrow 0 \\ & & \downarrow s & & \downarrow s' & & \downarrow id \\ 0 & \longrightarrow & E & \longrightarrow & Q & \longrightarrow & J \longrightarrow 0, \end{array}$$

where the left-hand square is a pushout. Since I, E are of type 1, s is RH -split. It is easy to show that an H -splitting of s induces an H -splitting of s' . Also, $\text{coker } s' \cong \text{coker } s$. Thus, in $a_R(G, H)$, we have

$$[P'] = [Q] - [\text{coker } s].$$

Now, $\text{coker } s$ is of type 1, so $[\text{coker } s] \in U$ by (3.22). Hence it suffices to show $[Q] \in U$. But J is irreducible of type 2; so, $J \cong X_n$. Thus, Q is an extension of X_n by E , and hence $Q \cong (X_n, E)_k$ for some k . Thus, $[Q] \in U$, and the proof of (3.23) is complete.

As we have noted before, (3.23) implies that $a_R(G, H)$ is generated by $[X_0], \dots, [X]_n, [(X_n, X_0)]$. Let us now show that these generators are independent. Suppose we have a relation

$$(3.25) \quad \sum_{i=0}^n x_i [X_i] + y \cdot [(X_n, X_0)] = 0$$

in $a_R(G, H)$, where x_i, y are integers. Then this relation also holds in the Grothendieck ring $K^0(RG) = a_R(G, 1)$. But $[(X_n, X_0)] = [X_n] + [X_0]$ in $K^0(RG)$. Since $K^0(RG)$ is free abelian on $[X_0], \dots, [X_n]$, it follows that $x_0 = -y = x_n$, $x_i = 0$ for $i = 1, \dots, n-1$. Mapping to $a_R(H)$ by the restriction map, we get $x_0[A] + x_n \cdot p^{n-1}[B] + y \cdot ([RH] + (p^{n-1} - 1)[B]) = 0$, using (3.14). Since $a_R(H)$ is free abelian with basis $[A], [B], [RH]$, it follows that $x_0 = y = x_n = 0$. Hence all coefficients in (3.25) are zero, and thus, the given generators are independent. Our proof shows also that the map

$$a_R(G, H) \rightarrow a_R(G, 1) \oplus a_R(H)$$

given by $[M] \mapsto ([M], [M_H])$ is a one-to-one mapping of rings. By [14] and [16], $a_R(G, 1) \oplus a_R(H)$ has no nonzero nilpotent elements. Hence we have shown

(3.26) **THEOREM.** *Let R be the ring of p -adic integers, let G be a cyclic group of order p^n and let H be the unique subgroup of order p in G . Then $a_R(G, H)$ is free as an additive group, with rank $n + 2$. As a ring, $a_R(G, H)$ has no nonzero nilpotent elements.*

Let us observe that (3.26) stands in contrast to the results in the modular case. Indeed, setting $\bar{R} = R/pR$, we have

(3.27) THEOREM. *The restriction map $\text{res}: a_R(G, H) \rightarrow a_R(H)$ is a ring isomorphism.*

Proof. See [10, Theorem 2.1].

Now, $a_R(H)$ is a free abelian group of rank p (see [14]). Thus the rank of $a_R(G, H)$ depends only on p . As we have just seen, the rank of $a_R(G, H)$ depends only on n . This is indeed a curious anomaly.

4. ELEMENTARY ABELIAN p -GROUPS

In this section, p is a prime, R is the ring of p -adic integers in the p -adic field K . Let G be an elementary abelian p -group of order p^n . Thus, G has generators a_1, \dots, a_n with relations $[a_i, a_j] = 1 = a_i^p$ for $i, j = 1, \dots, n$. Let $H = \langle a_1 \rangle$. We will calculate $a_R(G, H)$.

Denote by ζ a primitive p -th root of unity over K . From [1], we deduce that KG is the direct sum of the trivial G -module K , together with $(p^n - 1)/(p - 1)$ copies of $K[\zeta]$. The generators a_i act on $K[\zeta]$ as various powers of ζ ; thus RG acts through the discrete valuation ring $R[\zeta]$. As $R[\zeta]$ is a principal ideal domain, it follows that the nontrivial irreducible RG -lattices are all isomorphic to $R[\zeta]$, with the a_i acting as indicated above. Now, if k is an integer with $1 \leq k \leq p - 1$, then ζ^k is also a primitive p -th root of unity and $R[\zeta] = R[\zeta^k]$. Further, the matrix of the action of ζ on $R[\zeta]$ with respect to the R -basis $1, \zeta, \dots, \zeta^{p-2}$ is the same as that of the action of ζ^k with respect to the R -basis $1, \zeta^k, (\zeta^k)^2, \dots, (\zeta^k)^{p-1}$. Namely, both matrices are the companion matrix \mathbf{C} of the cyclotomic polynomial $\Phi_p(X)$ over K . Hence in choosing a full set of irreducible RG -lattices, we may assume that on each of them, a_1 acts either trivially or as multiplication by ζ . As in §3, we say that an RG -lattice X is of *type 1* if X is H -trivial, and that X is of *type 2* if $X_H \cong B^{(t)}$, some t .

Let X be an irreducible RG -lattice of type 2. Then X affords a matrix representation

$$\begin{aligned} a_1 &\rightarrow \mathbf{C} \\ a_k &\rightarrow \mathbf{C}^{n_k}, \quad k = 2, \dots, n, \end{aligned}$$

where the n_k are integers lying between zero and $p - 1$. Denote by T the trivial RG -lattice. We define an RG -lattice (X, T) by requiring that it afford the matrix representation

$$\begin{aligned} a_1 &\rightarrow \mathbf{D} \\ a_k &\rightarrow \mathbf{D}^{n_k}, \quad k = 2, \dots, n, \end{aligned}$$

where

$$D = \left(\begin{array}{c|cccc} 1 & 0 & 0 & \cdots & 1 \\ \hline 0 & & & & \\ 0 & & & & \\ \vdots & & & & \\ 0 & & & & \end{array} \begin{array}{c} \\ \\ \\ \\ \mathbf{C} \end{array} \right).$$

Then there are *RG*-exact sequences

$$(4.1) \quad 0 \longrightarrow X \xrightarrow{\mu} (X, T) \longrightarrow T \longrightarrow 0.$$

$$(4.2) \quad 0 \longrightarrow T \longrightarrow (X, T) \longrightarrow X \longrightarrow 0.$$

Neither sequence is *RH*-split, since $X_H \cong B$, $T_H \cong A$ and $(X, T)_H \cong RH$.

Imitating the proofs of (2.4) and (2.5), we may show

(4.3) LEMMA. Suppose that

$$0 \longrightarrow B \xrightarrow{\alpha} M \longrightarrow M/B \longrightarrow 0$$

is a nonsplit short exact sequence of *RH*-lattices. Then there is an *RH*-sublattice Y of M such that

- (i) $Y \cong RH$,
- (ii) $\alpha(B) \subseteq Y$,
- (iii) Y is an *RH*-summand of M_H .

(4.4) LEMMA. Suppose a diagram

$$\begin{array}{ccc} B & \xrightarrow{\alpha} & RH \\ \beta \downarrow & & \\ & & RH \end{array}$$

is given, where α, β are *RH*-monomorphisms with *R*-torsion-free cokernels. Then there is a commutative diagram

$$\begin{array}{ccc} B & \xrightarrow{\alpha} & RH \\ \beta \downarrow & \nearrow \mu & \\ & & RH, \\ & \nwarrow \theta & \end{array}$$

where μ, θ are *RH*-isomorphisms.

(4.5) LEMMA. *Let U be the subgroup of $a_R(G, H)$ generated by all $[X]$ such that X is irreducible and all $[(X, T)]$ such that X is irreducible of type 2. Then $U = a_R(G, H)$.*

Proof. Let M be an RG -lattice; we show $[M] \in U$ by induction on the R -rank of M . If M is irreducible or $M \cong (X, T)$ for some X , then $[M] \in U$. If all composition factors of M are of the same type, then in $a_R(G, H)$, $[M]$ equals the sum of its composition factors.

It remains to consider the case where M has composition factors of both types. Then there is an exact sequence

$$(4.6) \quad 0 \longrightarrow X \xrightarrow{\alpha} M \longrightarrow M/X \longrightarrow 0$$

with X irreducible of type 2. If (4.6) is RH -split, then $[M] = [X] + [M/X]$ in $a_R(G, H)$, whence $[M] \in U$ by the induction hypothesis. If (4.6) is not RH -split, then by (4.3), (4.4), and (3.17), we deduce

$$[M] = [(X, T)] - [T] + [M/X] \quad \text{in } a_R(G, H);$$

so, by the induction hypothesis, the proof is complete.

Now let X, Y be irreducible RG -lattices of type 2. Since $(X, T)_H \cong (Y, T)_H \cong RH$, and $T_H \cong A$, we may apply (2.5) and (3.16) to obtain $[(Y, T)] = [(X, T)] - [X] + [Y]$ in $a_R(G, H)$. Thus, we need only one of the $[(X, T)]$. Suppose that the irreducibles are

$$T, W_1, \dots, W_r, X_1, \dots, X_s,$$

where W_1, \dots, W_r are of type 1 and are not G -trivial and X_1, \dots, X_s are of type 2. Then, by (4.2.5) and the remarks above, we see that $a_R(G, H)$ is generated by $[T], [W_1], \dots, [W_r], [X_1], \dots, [X_s], [(X_1, T)]$. As in §3, we can show that these are independent. We obtain

(4.7) THEOREM. *Let R denote the p -adic integers, let G be an elementary abelian p -group and let H be a subgroup of order p . Then $a_R(G, H)$ is free abelian as an additive group, with rank $(p^n - 1)/(p - 1) + 2$. As a ring, $a_R(G, H)$ has no nonzero nilpotent elements.*

Let us remark that we have produced another counter example to the methods of the modular case. Indeed, Lam and Reiner [8, Theorem 5.1] have shown

(4.8) THEOREM. *Let G be a direct product $B \times K$ of finite groups B, K*

and let $H \subseteq K$. Assume that the field F of finite characteristic is a splitting field for B and all of its subgroups. Then, there is a ring isomorphism

$$a_F(B, 1) \otimes_{\mathbf{Z}} a_F(K, H) \cong a_F(G, H),$$

where \mathbf{Z} denotes the ring of rational integers.

Let $\bar{R} = R/pR$, $B = \langle a_2, \dots, a_n \rangle$, $H = K = \langle a_1 \rangle$. Then (4.8) yields a ring isomorphism

$$a_{\bar{R}}(B, 1) \otimes_{\mathbf{Z}} a_{\bar{R}}(H) \cong a_{\bar{R}}(G, H).$$

However, there is no ring isomorphism or even group isomorphism between $a_{\bar{R}}(B, 1) \otimes_{\mathbf{Z}} a_{\bar{R}}(H)$ and $a_{\bar{R}}(G, H)$ (since these groups have different ranks). Thus, (4.8) fails in the integral case. Finally, we note that (4.7) gives another counter example to the integral version of [10, Theorem 2.1].

5. QUATERNION AND DIHEDRAL GROUPS

Let R denote the ring of 2-adic integers. Let G be a nonabelian group of order 8, so G is quaternion or dihedral. Let $H = [G, G]$; then $H \triangleleft G$ and $|H| = 2$. By methods similar to those in §3 and §4, one can show

(5.1) THEOREM. $a_R(G, H) = U \oplus V$, where U is free abelian of rank six and $|V| \leq 2$.

The group V is generated by $[Y_1] - [Y_2]$, where Y_1, Y_2 are the two inequivalent RG -modules generating the unique irreducible KG -module of dimension > 1 . It seems quite difficult to determine whether or not $V = 0$.

On the basis of §3–§5, we conjecture that if $R = p$ -adic integers, G is a p -group, $H \triangleleft G$ and $|H| = p$, then $a_R(G, H)$ is free abelian of rank $e(G) + 1$, where $e(G)$ denotes the number of irreducible representations of G in the p -adic field. One may adapt the proof of (3.23) to give at least a partial proof of this conjecture.

6. MISCELLANY

An RG -lattice M is (G, H) -projective if every short exact sequence

$$0 \rightarrow N \rightarrow L \rightarrow M \rightarrow 0$$

of RG -lattices, which is RH -split is also RG -split. The subgroup $k_R(G, H)$ of $a_R(G)$ generated by all $[M]$ such that M is (G, H) -projective is an ideal, and there is a natural map

$$c: k_R(G, H) \rightarrow a_R(G, H)$$

given by $c([M]) = [M]$. c is the *Cartan homomorphism*. By the same proof as in [8], one may show

(6.1) THEOREM. *Let R be a complete discrete valuation ring. Then c is monic.*

From this, one can deduce

(6.2) COROLLARY. *Let R be a Dedekind ring such that the Jordan–Zassenhaus Theorem holds for RG -lattices and the residue fields of R have finite characteristics. Then $\ker c$ is finite.*

(6.3) COROLLARY. *Let R be a discrete valuation ring (not necessarily complete) with finite residue field. Then c is monic.*

(6.4) COROLLARY. *Let R be a complete discrete valuation ring and assume that the Jordan–Zassenhaus Theorem holds for RH -lattices. Suppose that $a_R(G, H)$ is finitely generated. Then the number $n(RH)$ of isomorphism classes of indecomposable RH -lattices is finite.*

(6.5) COROLLARY. *Let K be an algebraic number field which is a splitting field for G , and let R be the ring of algebraic integers in K . Suppose that $a_R(G, H)$ is finitely generated. Then $n(RH) < \infty$.*

On the basis of (6.4), (6.5), and the results of [9], we conjecture that $a_R(G, H)$ is finitely generated if and only if $n(RH) < \infty$.

One may also show that two other modular results do not carry over to the integral case. Namely, Lam and Reiner [6] and Dress [4] have shown that the cokernel of c is a torsion group in the modular case. In contrast, we have

(6.6) PROPOSITION. *Let $R = p$ -adic integers, G a p -group, H a subgroup of order p . If $H \neq G$, then the cokernel of c has elements of infinite order.*

Also, in the modular case, it is shown in [10] that the restriction map

$$\text{res}: a_F(G, H) \rightarrow a_F(H)$$

is often monic or even an isomorphism. We have

(6.7) PROPOSITION. *Let R be a domain of characteristic zero. Let $H \neq G$. Then $\text{res}: a_R(G, H) \rightarrow a_R(H)$ is not monic.*

REFERENCES

1. R. AYOUB AND C. AYOUB, On the group ring of a finite abelian group, *Bull. Austral. Math. Soc.* **1** (1969), 245–262.
2. S. B. CONLON, Decompositions induced from the Burnside algebra, *J. Algebra* **10** (1968), 102–122.
3. S. B. CONLON, Modular representations of $C_2 \times C_2$, *J. Austral. Math. Soc.* **10** (1969), 363–366.
4. A. DRESS, On relative Grothendieck rings, *Bull. Amer. Math. Soc.* **75** (1969), 955–958.
5. A. HELLER AND I. REINER, Representations of cyclic groups in rings of integers I, *Ann. of Math. (2)* **76** (1962), 73–92.
6. T.-Y. LAM AND I. REINER, Relative Grothendieck groups, *J. Algebra* **11** (1969), 213–242.
7. T.-Y. LAM AND I. REINER, Relative Grothendieck rings, *Bull. Amer. Math. Soc.* **75** (1969), 496–498.
8. T.-Y. LAM AND I. REINER, Reduction theorems for relative Grothendieck rings, *Trans. Amer. Math. Soc.* **142** (1969), 421–435.
9. T.-Y. LAM AND I. REINER, Finite generation of Grothendieck rings relative to cyclic subgroups, *Proc. Amer. Math. Soc.* **23** (1969), 481–489.
10. T.-Y. LAM AND I. REINER, Restriction maps on relative Grothendieck rings, *J. Algebra* **14** (1970), 260–298.
11. T.-Y. LAM AND I. REINER, An excision theorem for Grothendieck rings, *Math. Z.* **115** (1970), 153–164.
12. T.-Y. LAM, I. REINER, AND D. WIGNER, Restrictions of representations over fields of characteristic p , *Proc. Symp. Pure Math. (Amer. Math. Soc.)* **21** (1971) 99–106.
13. S. MACLANE, “Homology,” Springer-Verlag, Berlin, 1963.
14. I. REINER, Integral representation algebras, *Trans. Amer. Math. Soc.* **124** (1966), 111–121.
15. R. SWAN, Induced representations and projective modules, *Ann. of Math. (2)* **71** (1960), 552–578.
16. R. SWAN, The Grothendieck ring of a finite group, *Topology* **2** (1963), 85–110.